

การประเมินการแลกเปลี่ยนก๊าซคาร์บอนไดออกไซด์ของอ้อยตอหนึ่ง โดยวิธี Eddy Covariance Technique

Evaluation of Carbon Dioxide Flux of the First Ratoon Cane by Eddy Covariance Technique

นงภัทร ไชยชนะ¹ ทิวา พาโคทอม¹ เจษฎา ภัทรเลอพงศ์² และ เชษฐ สาทริกจิ³

Nongpat Chaichana¹, Tiwa Pakoktom¹, Jessada Phattaralerphong² and Jate Sathornkich³

บทคัดย่อ

การทดลองครั้งนี้เป็นการประเมินการแลกเปลี่ยนก๊าซคาร์บอนไดออกไซด์ (CO₂) ของอ้อยตอ 1 ณ ศูนย์ส่งเสริมอุตสาหกรรมอ้อยและน้ำตาลทรายภาคกลาง จังหวัดกาญจนบุรี ตั้งแต่เดือนมิถุนายน 2553 ถึงเดือนเมษายน 2554 โดยเทคนิค Eddy Covariance จากข้อมูลการแลกเปลี่ยน CO₂ ของอ้อยตอ 1 ตลอดการทดลอง 11 เดือน พบว่าค่าการแลกเปลี่ยน CO₂ กับบรรยากาศมีค่าอยู่ระหว่าง -2.4-2.8 mgCO₂m⁻²s⁻¹ โดยค่าของ CO₂ มีค่าเป็นลบในช่วงเวลากลางวัน แสดงถึงพืชดูดซับ CO₂ โดยกระบวนการสังเคราะห์ด้วยแสง แต่ค่าของ CO₂ มีค่าเป็นบวกในช่วงเวลากลางวัน แสดงถึง CO₂ ถูกปลดปล่อยออกจากแปลงโดยกระบวนการหายใจของพืชและดิน เมื่อเปรียบเทียบปริมาณ CO₂ ที่แปลงอ้อยตอ 1 สามารถดูดซับได้ในแต่ละระยะการเจริญเติบโต พบว่าปริมาณ CO₂ ที่ถูกดูดซับในแต่ละระยะการเจริญเติบโตมีค่าที่แตกต่างกัน โดยระยะออกสามารถดูดซับ CO₂ ได้น้อยที่สุด คือ 63.1 gCO₂m⁻² ส่วนระยะย่างปล้องสามารถดูดซับ CO₂ ได้มากที่สุด คือ 2,448.3 gCO₂m⁻² ทั้งนี้ปริมาณ CO₂ ที่แปลงอ้อยตอ 1 สามารถดูดซับได้ทั้งหมดตลอดฤดูปลูกมีค่าเท่ากับ 4,300.4 gCO₂m⁻² จากการทดลองครั้งนี้สามารถสรุปได้ว่า แปลงอ้อยตอ 1 เป็นแหล่งปลดปล่อย CO₂ ในเวลากลางคืน ส่วนในช่วงเวลากลางวันจะเป็นแหล่งดูดซับ CO₂ และตลอดฤดูปลูกอ้อยตอ 1 เป็นแหล่งดูดซับ CO₂ จากบรรยากาศ

ABSTRACT

The propose of this study was to evaluate CO₂ flux and accumulative CO₂ absorbance of the first ratoon cane by using Eddy Covariance Technique. First ratoon cane was produced at Cane and Sugar Industry Promotion Center, Kanchanaburi province, from June 2010 to April 2011. From the results, the CO₂ flux was fluctuated from -2.4-2.8 mg CO₂m⁻²s⁻¹. The negative values showed in the day time and it means CO₂ was absorbed by plant via photosynthesis. On the other hand the positive values means CO₂ was emitted by plant and soil respiration in night time. Different CO₂ flux was observed in each growth stages.

¹คณะเกษตร กำแพงแสน มหาวิทยาลัยเกษตรศาสตร์ วิทยาเขตกำแพงแสน จ.นครปฐม 73140

¹Faculty of Agriculture at Kamphaeng Saen, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom 73140

²คณะทรัพยากรธรรมชาติและอุตสาหกรรมเกษตร มหาวิทยาลัยเกษตรศาสตร์ วิทยาเขตเฉลิมพระเกียรติ จังหวัดสกลนคร จ.สกลนคร 47000

²Faculty of Natural Resource and Agro-Industry, Kasetsart University, Chalermprakiat Sakon Nakorn, Sakon Nakorn, 47000

³ภาควิชาพืชสวน คณะเกษตร มหาวิทยาลัยเกษตรศาสตร์ เขตจตุจักร กรุงเทพฯ 10900

³Department of Horticulture, Faculty of Agriculture, Kasetsart University, Chatuchak, Bangkok 10900

CO₂ flux was lower at germination stage (June 2010) and the total CO₂ absorbance at this stage was 63.1 g CO₂m⁻². CO₂ flux reached its maximum at stalk elongation stage (September 2010-January 2011) and total CO₂ absorbance at this stage was 2,448.3 g CO₂m⁻². Totally, CO₂ absorbance for the entire period of experiment was 4,300.4 g CO₂m⁻². From the results, it was concluded that the 1st ratoon sugarcane ecosystem was a CO₂ source during night hours and a CO₂ sink during the day time. Finally, for the entire growing season the 1st ratoon cane system is the sink of CO₂.

Key Words: CO₂ flux, Eddy Covariance Technique, first ratoon cane

e-mail address: nongpat.chai@gmail.com

INTRODUCTION

Sugarcane is an important economic crop in Thailand and the world not only for sugar production but increasingly also for a bioenergy crop due to its phenomenal dry matter production capacity (Marcelo *et al.*, 2007). Global sugarcane production now is estimated at 1,250 million tons a year while the total production of sugarcane in Thailand is about 106 million tons per year. Total planting area of sugarcane in Thailand is 1.42 million ha (Office of the Cane and Sugar Board, 2012). Carbon dioxide (CO₂) exchange between terrestrial ecosystem and the atmosphere one of the key process that affects atmospheric CO₂ concentration. In order to assess the role of the terrestrial ecosystem in the global CO₂ budget at present, and to predict its changes in the future under global warming, long-term observation of CO₂ exchange has been done in various ecosystems in the world. These observation sites are mostly located in forest ecosystems because these are believed to be the most influential terrestrial ecosystem in the global CO₂ budget. However, non-forest ecosystem, such as grasslands, wetland or agricultural field also cover some part of the land area and contribute to regional and global CO₂ budgets (Saito *et al.*, 2005). Gas fluxes can be measured using many methods and Eddy Covariance Technique (EC) is the new method which does not interfere with process of gas exchange between the surface source and the atmosphere (Kuo-Hsin *et al.*, 2010). This technique has been used to measure CO₂ flux, and many studies have been published (Schrier-Uijl *et al.*, 2010). However, most of gas exchange studies involved short-term measurements. In this paper, we present seasonal variation of CO₂ flux at a customarily cultivated single cropping sugarcane crop base on the results of a single season-long measurement of CO₂ flux using the EC technique.

MATERIALS AND METHODS

1. Site description and period of measurement

This site was conducted at the first ratoon cane which is located at Cane and Sugar Industry Promotion Center Region 1, Kanchanaburi province, in Western Thailand. The site is located at latitude

14.03°N and longitude 99.68°E with an elevation of 22.37 m above mean sea level. The soil of the sugarcane field is sandy loam. A CO₂ flux measurement and meteorological tower was 6.0 m high and was erected in the center of plot area. The sugarcane fields around the tower were managed as single sugarcane-cropping fields following by a common management in this area. In this study, the data of the 1st ratoon cane growing season (9 June 2010-31 April 2011) were analyzed.

2. Meteorological data measurement

The meteorological parameter consisted of solar radiation (Rs), net radiation (Rn), total amount of rain, wind speed and wind direction, air temperature and relative humidity. All of the meteorological instruments were installed on the tower at a height of 6.0 m. Soil moisture contents which measured by EC-20 (Decagon Devices, Inc.) were buried at depths of 15 and 45 cm in the soil around the tower. The meteorological data were collected and averaged every 30 minutes and stored in data logger CR1000 (Campbell Scientific, Inc.).

3. Plant growth

Plant-cane was planted in 30 June 2009 and the first ratoon cane started from 9 June 2010. Growth characteristics consisted of plant height, stem diameter, fresh and dry weight, leaf area (LA) and leaf area index (LAI). The samples were taken from 5 plots around the tower at monthly intervals commencing from 60 days after plant-cane was harvested until the next harvest. Each samples consisted of the above-ground portions of all shoots, the shoots were separated into stem, leaf blade and leaf sheath. When the amount of sample was in excess, a portion of representative subsample was used for dry weight determination, extrapolated for a whole sample, which was designated as the total dry matter. All samples were oven-dried at 80°C until a constant weight was reached.

4. Eddy Covariance flux measurement

CO₂ flux was measured by the Eddy Covariance technique. Three components of wind velocity and temperature fluctuation were measured with a sonic anemometer (CSAT-3; LI-COR, Inc., Lincoln, NE, USA). The densities of CO₂ was measured with an open-path infrared gas analyzer (IRGA) (LI-7500; LI-COR, Inc., Lincoln, NE, USA). The sensor heads of the sonic anemometer and IRGA were mounted the tower at a height of 2.0 m above the plant canopy. The data from the sonic anemometer and IRGA were sampled at 20 Hz using a 16-bit digital data recorder and stored in data logger CR3000. Half-hourly flux density of the CO₂ was calculated from the covariance between the vertical wind velocity and the respective quantities. All data were downloaded to computer every week and the quality of data were checked against the standard meteorology (Makoto *et al.*, 2005). The CO₂ flux data were shown in plus (+) and minus (-), the meaning of plus is CO₂ was released from ecosystem by plant and soil respirations to atmosphere, and minus indicated CO₂ was absorbed into crop community through photosynthesis.

RESULTS

Meteorological data

Meteorological condition in the first ratoon cane during 9 June 2010-30 April 2011 was shown in Figure 1. Daily average solar radiation (Rs) variation ranged from 3.29-27.56 $\text{MJm}^{-2}\text{d}^{-1}$, lower levels of solar radiation in March 2011. Net radiation (Rn) was increased after the sunrise and reached its maximum at noon. The daily average of Rn variation ranged from 0.63-22.55 $\text{MJm}^{-2}\text{d}^{-1}$. The trend of average air temperature (Ta) and average relative humidity (RH) were shown in Figure 2. The time series of average air temperature (Ta) fluctuation was between 17.9-31.1°C, its minimum was 17.9°C in March 2011 and its maximum was 31.1°C in April 2011. The average relative humidity (RH) was 71% and its minimum was 53%. Total amount of rain fall during the study period was 345 mm and its maximum was 61.1 mm per day which occurred in 29 June 2010 (Figure 3). Soil moisture content from soil dept 10 and 20 cm varied in the range of 10-34% and 10-40%V/V, respectively (Figure 3).




Figure 1 Time series of solar radiation (Rs) and net radiation (Rn) at the experimental site from 9 June 2010-30 April 2011




Figure 2 Time series of average air temperature (Ta) and average relative humidity (RH) at the experimental site from 9 June 2010-30 April 2011




Figure 3 Total amount of rain and soil moisture content at the experimental site during 9 from 2010-30 April 2011

Plant growth parameters

Plant height and diameter increased gradually from the beginning of measurement and reached the saturated point at age 7 month. Plant height was reached a maximum of 152 cm at maturity stage. The LAI showed a gradual increase from the tillering stage and reached a maximum at stalk elongation stage (Figure 4)




Figure 4 The plant height (cm) and leaf area index (LAI) of the first ratoon cane

Daily trend of CO₂ flux

All of the data set of the first ratoon cane were analyzed for the daily trend of CO₂ flux. The daily trend of CO₂ flux was presented in Figure 5. Throughout the measurement period, the daily values ranged from $-0.5 \text{ mg CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ to $0.13 \text{ mg CO}_2 \text{ m}^{-2} \text{ s}^{-1}$. It can be inferred that the fluxes of CO₂ were always positive during night hours with an average night time value of $0.11 \text{ mg CO}_2 \text{ m}^{-2} \text{ s}^{-1}$, whereas during the day time the flux was negative with a corresponding value of $-0.17 \text{ mg CO}_2 \text{ m}^{-2} \text{ s}^{-1}$. Total CO₂ released of $4.8 \text{ g CO}_2 \text{ m}^{-2}$ and total CO₂ absorbed of $12.4 \text{ g CO}_2 \text{ m}^{-2}$ were observed.




Figure 5 Daily trend of CO₂ flux of the first ratoon cane from 9 June 2010-30 April 2011

Seasonal trend of CO₂ flux

Seasonal variation of CO₂ flux was shown on Figure 6. The daily values ranged between 0.05 to 31.1 g CO₂m⁻²d⁻¹. The CO₂ flux summation at germination stage (0-1 month), tillering stage (2-3 month), stalk elongation stage (4-8 month) and maturity stage (9-11 month) were 63.1, 838.2, 2,448.3 and 950.8 gCO₂m⁻², respectively. At stalk elongation stage the CO₂ flux reached maximum cause of high LAI and this period which are 5 months in length. Totally, CO₂ absorbance of 4,300.4 gCO₂m⁻² of the 1st ratoon cane was recorded.




Figure 6 Seasonal trend of CO₂ flux of the first ratoon cane from 9 June 2010-30 April 2011

DISCUSSION

During the day time plant photosynthesis leads to uptake of CO₂ from both the atmosphere and from respired CO₂ emitted by the soil. Respiration at night leads to an efflux of CO₂ to the atmosphere (Miyata *et al.*, 2000). Watcharapirak (2009) found that the carbon storage content of sugarcane plantation in one crop yield consists of the carbon storage in sugarcanes and the organic carbon deposits in the soil. This carbon storage content was increases when the sugarcane grows. The results were similar to the reported in paddy fields by Pakoktom *et al.* (2009) that CO₂ flux was lower in stage I because of small LAI, then, gradually in increased and reached its maximum in the later stage. Moreover, CO₂ flux decreased in stage IV, probably due to the leaf senescence. Net exchange of CO₂ between crop community and the atmosphere is controlled by several biological and physical processes. It is well accepted that CO₂ storage with in the forest canopy has to be taken into account wherever we discuss the net CO₂ exchange between the ecosystem and the atmosphere (Aubinet *et al.*, 2001).

CONCLUSION

During the night time the first ratoon cane was a CO₂ source originating from plant and soil respiration and during the day time became a CO₂ sink during the day time as plant photosynthesis advanced. However, for 24 hr period, it was a CO₂ sink. From this result it was concluded that the 1st ratoon sugarcane ecosystem is a sink of CO₂.

ACKNOWLEDGEMENTS

This study was financially supported by the Office of the Cane and Sugar Board. Many thanks are due to DORAS center for their encouragement and field instrumentations. Finally, we are grateful to the Cane and Sugar Industry Promotion Center Region 1 for providing the experimental plots and field assistance.

LITERATURE CITED

- Aubinet, M., B. Chermanne, M. Vandenhaute, B. Longdoz, M. Yernaux and E. Laitat. 2001. Long term carbon dioxide exchange above a mixed forest in the Belgian Ardennes. **Agricultural and Forest Meteorology** 108: 293-315.
- Kuo-Hsin, T., J. Tsai, A. Alagesan, B. Tsuang, M. Yao and P. Kuo. 2010. Determination of methane and carbondioxide fluxes during the rice maturity period in Taiwan by combining profile and eddy covariance measurements. **Agricultural and Forest Meteorology** 150: 825-859.

- Makoto, S, A. Miyata, H. Nagai and T. Yamada. 2005. Seasonal variation of carbon dioxide exchange in rice paddy field in Japan. **Agricultural and Forest Meteorology** 135: 93-109.
- Marcelo de A. S., J. L. Jifon, J. A. G. Da Silva and V. Sharma. 2007. Use of physiological parameters as fast tools to screen for drought tolerance in sugarcane. **Braz. J. Plant Physiol.** 19(3): 193-201.
- Miyata, A., R. Leuning, O. T. Demeand, J. Kim and Y. Harazono. 2000. Carbon dioxide and methane fluxes from an intermittently flooded paddy field. **Agricultural and Forest Meteorology** 102: 287-303.
- Office of the Cane and Sugar Board 2012. Report on Cane and Sugar Production 2011/12. Available Source: <http://www.ocsb.go.th/upload/journal/fileupload/923-3963.pdf>, 14 September, 2012.
- Pakoktom, T., M. Aoki, P. Kasemsap, S. Boonyawat and P. Attarod. 2009. CO₂ and H₂O fluxes ratio in paddy fields of Thailand and Japan. **Hydrological Research Letter** 3: 10-13.
- Saito, M., A. Miyata, H. Nagai and T. Yamada. 2005. Seasonal variation of carbon dioxide exchange in rice paddy field in Japan. **Agricultural and Forest Meteorology** 135: 93-109.
- Schrier-Uijl, A.P., P.S. Kroon, A. Hensen, P.A. Leffelaar, F. Berendse and E.M. Veenendaal. 2010. Comparison of chamber and eddy covariance-based CO₂ and CH₄ emission estimates in a heterogeneous grass ecosystem on peat. **Agricultural and Forest Meteorology** 150: 825-831.
- Watcharapirak, W. 2009. The estimation of carbon storage in various growth stages of sugarcane in Sisatchanalai district, Sukhothai province, Thailand. Master Thesis, Mahidol University.